

Spins of primordial black holes formed in the matter-dominated era

Tomohiro Harada (Rikkyo U)


13/12/2017, COSPA2017 @ YITP

This talk is based on


- Harada, Yoo (Nagoya U), Kohri (KEK), Nakao (OCU) & Jhingan (YGU), 1609.01588
- Harada, Yoo, Kohri, & Nakao, 1707.03595

Primordial black hole (PBH)

- PBH = Black hole formed in the early Universe
 - Probe into the early Universe, high-energy physics, and quantum gravity. May act as γ -ray or X-ray sources, dark matter candidate, and gravitational wave sources. (e.g. Carr et al. (2010), Carr et al. (2016))
 - LIGO BBHs may be made of PBHs. (Sasaki et al. (2016), Bird et al. (2016), Clesse & Garcia-Bellido (2017))
 - The observation of BH spins has attracted great attention. (e.g. McClintock (2011), Abbott et al. (2017))



(a) Carr et al. (2016)




(b) LIGO Collaboration (2017)

PBH formation in the matter-dominated (MD) era

- Pioneered by Khlopov & Polnarev (1980). Recently motivated by early MD phase scenarios such as inflaton oscillations, phase transitions, and superheavy metastable particles.
- Primordial perturbations may collapse to PBHs. If pressure is negligible, nonspherical effects play crucial roles.
 - The triaxial collapse of dust leads to a “pancake” singularity. (Lin, Mestel & Shu 1965, Zeldovich 1969)



- The effect of angular momentum may halt gravitational collapse or spin the formed PBHs.



- We here rely on the Newtonian approximation to deal with complicated nonspherical dynamics analytically.

Zeldovich approximation

- Zeldovich approximation (ZA) (1969)
Extrapolate the Lagrangian perturbation theory in the linear order in Newtonian gravity to the nonlinear regime.

$$\mathbf{r}_i = a(t)\mathbf{q}_i + \mathbf{b}(t)\mathbf{p}_i(\mathbf{q}_j),$$

where $\mathbf{b}(t) \propto a^2(t)$ denotes a linear growing mode.


- We can take the coordinates in which

$$\frac{\partial \mathbf{p}_i}{\partial \mathbf{q}_j} = \text{diag}(-\alpha, -\beta, -\gamma),$$

where we can assume $\infty > \alpha \geq \beta \geq \gamma > -\infty$.

- We assume that α , β and γ are constant over the smoothing scale.
- We normalise \mathbf{b} so that $(\mathbf{b}/a)(t_i) = \mathbf{1}$ at horizon entry $t = t_i$.

Application of the hoop conjecture to the pancake collapse



- Hoop conjecture (Thorne 1972): The collapse results in a BH if and only if $C \lesssim 4\pi GM/c^2$, where C is the circumference of the pancake singularity.
- Then, we obtain a BH criterion:


$$h(\alpha, \beta, \gamma) := \frac{C}{4\pi Gm/c^2} = \frac{2}{\pi} \frac{\alpha - \gamma}{\alpha^2} E \left(\sqrt{1 - \left(\frac{\alpha - \beta}{\alpha - \gamma} \right)^2} \right) \lesssim 1,$$

where $E(e)$ is the complete elliptic integral of the second kind.

- If $h \gtrsim 1$? : It does not immediately collapse to a BH.

Spin angular momentum within the region to collapse

- Region V : to collapse in the future



- Angular momentum within V with respect to the COM in the Eulerian coordinates

$$\mathbf{L} = \rho_0 a^4 \left(\int_V \mathbf{x} \times \mathbf{u} d^3\mathbf{x} + \int_V \mathbf{x} \delta \times \mathbf{u} d^3\mathbf{x} - \frac{1}{V} \int_V \mathbf{x} \delta d^3\mathbf{x} \times \int_V \mathbf{u} d^3\mathbf{x} \right),$$

where $\mathbf{x} := \mathbf{r}/a$, $\mathbf{u} := aD\mathbf{x}/Dt$, $\delta := (\rho - \rho_0)/\rho_0$, and $\psi := \Psi - \Psi_0$.

- Linearly growing mode of perturbation

$$\delta_1 = \sum_{\mathbf{k}} \hat{\delta}_{1,\mathbf{k}}(t) e^{i\mathbf{k}\cdot\mathbf{x}}, \quad \psi_1 = \sum_{\mathbf{k}} \hat{\psi}_{1,\mathbf{k}}(t) e^{i\mathbf{k}\cdot\mathbf{x}}, \quad \mathbf{u}_1 = \sum_{\mathbf{k}} \hat{\mathbf{u}}_{1,\mathbf{k}}(t) e^{i\mathbf{k}\cdot\mathbf{x}},$$

where

$$\hat{\delta}_{1,\mathbf{k}} = A_{\mathbf{k}} t^{2/3}, \quad \hat{\psi}_{1,\mathbf{k}} = -\frac{2}{3} \frac{a_0^2}{k^2} A_{\mathbf{k}}, \quad \hat{\mathbf{u}}_{1,\mathbf{k}} = i a_0 \frac{\mathbf{k}}{k^2} \frac{2}{3} A_{\mathbf{k}} t^{1/3}.$$

1st-order effect

$$\mathbf{L} = \rho_0 a^4 \left(\int_V \mathbf{x} \times \mathbf{u} d^3\mathbf{x} + \int_V \mathbf{x} \delta \times \mathbf{u} d^3\mathbf{x} - \frac{1}{V} \int_V \mathbf{x} \delta d^3\mathbf{x} \times \int_V \mathbf{u} d^3\mathbf{x} \right)$$

- If ∂V is not a sphere, the 1st term contribution grows as $\propto \mathbf{a} \cdot \mathbf{u} \propto t$.
- If we assume V is a triaxial ellipsoid with axes (A_1, A_2, A_3) , we find

$$\langle \mathbf{L}_{(1)}^2 \rangle^{1/2} \simeq \frac{2}{5\sqrt{15}} q \frac{MR^2}{t} \langle \delta^2 \rangle^{1/2},$$

where $r_0 := (A_1 A_2 A_3)^{1/3}$, $R := a(t)r_0$ and

$q := \sqrt{\frac{Q_{ij}Q_{ij}}{3(\frac{1}{5}Mr_0^2)^2}}$ is a nondimensional

reduced quadrupole moment of V . (Cf.

Catelan & Theuns 1996)




Figure: The 1st-order effect can grow if ∂V is not a sphere.

2nd-order effect

$$\mathbf{L} = \rho_0 a^4 \left(\int_V \mathbf{x} \times \mathbf{u} d^3\mathbf{x} + \int_V \mathbf{x} \delta \times \mathbf{u} d^3\mathbf{x} - \frac{1}{V} \int_V \mathbf{x} \delta d^3\mathbf{x} \times \int_V \mathbf{u} d^3\mathbf{x} \right)$$

- Even if ∂V is a sphere, the remaining contribution grows as 1st order \times 1st order $\propto \mathbf{a} \cdot \delta \cdot \mathbf{u} \propto t^{5/3}$.

$$\langle \mathbf{L}_{(2)}^2 \rangle^{1/2} = \frac{2}{15} I \frac{MR^2}{t} \langle \delta^2 \rangle,$$

where δ hereafter is the density perturbation averaged over V . $R := a(t)r_0$. We assume $I = O(1)$. (Cf. Peebles 1969)




Figure: The 2nd-order effect can grow due to the mode coupling.

The application of the Kerr bound to the PBH formation

- Time evolution of V and angular momentum

- Horizon entry ($t = t_H$): $a_{r_0} = cH^{-1}$, $\delta_H := \delta(t_H)$, $\sigma_H := \langle \delta^2 \rangle^{1/2}$
- Maximum expansion ($t = t_m$): $\delta(t_m) = 1$, typically $t_m = t_H \sigma_H^{-3/2}$
- $a_* := L/(GM^2/c)$ at $t = t_m$

$$\langle a_{*(1)}^2 \rangle^{1/2} = \frac{2}{5} \sqrt{\frac{3}{5}} q \sigma_H^{-1/2}, \langle a_{*(2)}^2 \rangle^{1/2} = \frac{2}{5} I \sigma_H^{-1/2}, a_* \simeq \max(\langle a_{*(1)}^2 \rangle, \langle a_{*(2)}^2 \rangle)$$

- For $t > t_m$, the evolution of V decouples from the cosmological expansion and hence a_* is kept almost constant.

- Consequences

- Supercritical angular momentum: typically $\langle a_*^2 \rangle^{1/2} \gtrsim 1$ if $\sigma_H \lesssim 0.1$
- Most of the PBHs have $a_* \simeq 1$. This contrasts with small spins ($a_* \lesssim 0.4$) of PBHs formed in the RD era. (Chiba & Yokoyama (2017))
- Suppression: The Kerr bound implies that a_* is typically too large for direct collapse to a BH.

Spin distribution

- Spin distribution of PBHs formed in the MD era




Figure: The distribution function normalised by the peak value. We assume a Gaussian distribution for the density perturbation. Each curve is labelled with the value of σ_H .

- The region with smaller δ_H has larger a_* . This implies that there appears a threshold δ_{th} below which the angular momentum halts the collapse to a black hole due to the Kerr bound.

Numerical calculation of PBH production rate

- Triple integral for β_0 ($\theta(x)$ is a step function.)

$$\beta_0 \simeq \int_0^\infty d\alpha \int_{-\infty}^\alpha d\beta \int_{-\infty}^\beta d\gamma \theta[\delta_H(\alpha, \beta, \gamma) - \delta_{\text{th}}] \theta[1 - h(\alpha, \beta, \gamma)] w(\alpha, \beta, \gamma),$$

where we use $w(\alpha, \beta, \gamma)$ given by Doroshkevich (1970).




Figure: The red lines are due to both angular momentum and anisotropy. The 1st-order effect depends on q . The black solid line is solely due to anisotropy.

- We have also derived semianalytic formulae for β_0 .


Summary

- PBHs may form not only in the RD era but also in the (early) MD era by primordial cosmological fluctuations.
- In the MD era, the effect of anisotropy gives $\beta_0 \simeq 0.05556\sigma_H^5$, while the effect of angular momentum gives further suppression for the smaller values of σ_H .
- PBHs formed in the MD era mostly have large spins ($a_* \simeq 1$) in contrast to the small spins ($a_* \lesssim 0.4$) of PBHs formed in the RD era.

Anisotropic collapse in the ZA


- The triaxial ellipsoid of a Lagrangian ball (assumption)

$$\begin{cases} r_1 = (a - \alpha b)q \\ r_2 = (a - \beta b)q \\ r_3 = (a - \gamma b)q \end{cases}$$



- Evolution of the collapsing region:

- Horizon entry ($t = t_i$): $a(t_i)q = cH^{-1}(t_i) = r_g := 2Gm/c^2$.
- Maximum expansion ($t = t_f$): $\dot{r}_1(t_f) = 0$ giving $r_f := r_1(t_f) = r_g/(4\alpha)$.
- Pancake singularity ($t = t_c$): $r_1(t_c) = 0$ giving $a(t_c)q = 4r_f = r_g/\alpha$.



Application of the Kerr bound to the rotating collapse

- Technical assumption

$$|\mathbf{L}_{(1)}| \simeq \frac{2}{5\sqrt{15}} q \frac{MR^2}{t} \delta, \quad |\mathbf{L}_{(2)}| \simeq \frac{2}{15} I \frac{MR^2}{t} \langle \delta^2 \rangle^{1/2} \delta.$$

- The above assumption implies

$$a_{*(1)} = \frac{2}{5} \sqrt{\frac{3}{5}} q \delta_H^{-1/2}, \quad a_{*(2)} = \frac{2}{5} I \sigma_H \delta_H^{-3/2}, \quad a_* = \max(a_{*(1)}, a_{*(2)}).$$

- The Kerr bound $a_* \leq 1$ gives a threshold δ_{th} for δ_H , where

$$\delta_{\text{th}} = \max(\delta_{\text{th}(1)}, \delta_{\text{th}(2)}), \quad \delta_{\text{th}(1)} := \frac{3 \cdot 2^2}{5^3} q^2, \quad \delta_{\text{th}(2)} := \left(\frac{2}{5} I \sigma_H \right)^{2/3}.$$

Discussion of PBH production

- Semianalytic estimate (black dashed line and blue dashed line)

$$\beta_0 \simeq \left\{ \begin{array}{ll} 2 \times 10^{-6} f_q(q_c) I^6 \sigma_H^2 \exp \left[-0.15 \frac{I^{4/3}}{\sigma_H^{2/3}} \right] & \text{(2nd-order effect)} \\ 3 \times 10^{-14} \frac{q^{18}}{\sigma_H^4} \exp \left[-0.0046 \frac{q^4}{\sigma_H^2} \right] & \text{(1st-order effect)} \\ 0.05556 \sigma_H^5 & \text{(anisotropic effect)} \end{array} \right. ,$$

where $f_q(q_c)$: the ratio of regions with $q < q_c = O(\sigma_H^{1/3})$.

- σ_H in terms of P_ζ :

$$\sigma_H^2 \simeq \left(\frac{2}{5} \right)^2 P_\zeta(k_{BH}).$$