


# Hawking radiation from supertranslated horizon

Yoji Koyama  
(NCTS, Tawian)

ongoing work with Chong-Sun Chu (NCTS/National Tsing Hua U)

# Introduction

- BMS supertranslations - asymptotic symmetries of asymptotically Minkowski spacetime
- Supertranslation hair on black hole - new kind of hair which may be used to distinguish black holes with the same  $M, Q, J$   
[Hawking, Perry and Strominger (2016)]
- Question: Does Hawking radiation carry some information of supertranslation hair?  A: Yes, for dynamical Vaidya BH
- Hawking radiation as tunneling [Parikh and Wilczek]

## □ BMS supertranslation

[Bondi, Metzner, van der Burg, Sachs]

Diffeomorphism which preserves Bondi gauge conditions and asymptotic flatness conditions at null infinity

- Vector field  $\zeta$  which generates BMS supertranslation is

$$\zeta_f = f\partial_v - \frac{1}{2}D^2 f\partial_r + \frac{1}{r}D^A f\partial_A, \quad f = f(\Theta) : \text{arbitrary function of angle}$$

Advanced Bondi coordinates  $(v, r, z, \bar{z})$

- Supertranslations are parametrized by a function  $\Rightarrow$  infinitely many

\* Charge associated with supertranslation:  $Q_f = \int_{i^0} f(\Theta) m d^2\Omega$

$$f(\Theta) = \sum_{l,m} a_{lm} Y_{lm}(\Theta) \quad m : \text{Bondi mass}$$

## □ Supertranslation hair

Schwarzschild spacetimes with different supertranslations

$$(M_{\text{ADM}}, Q_{f(\Theta)}) \quad (\tilde{M}_{\text{ADM}}, Q_{\tilde{f}(\Theta)})$$

- Classically,  $M_{\text{ADM}} = \tilde{M}_{\text{ADM}}$ ,  $Q_f = Q_{\tilde{f}}$        $\{Q_f, Q_{f'}\}_D = 0$  [Barnich et al.]

(Superrotation charge can be interpreted as the classical hair.)  
[Hawking et al.]

- Quantum mechanically, it is interesting to see if Hawking radiation depends on  $f$ . If so, black holes with different supertranslations can be distinguished

## □ Our setup

- We consider dynamical black hole as black hole will be dynamical due to the backreaction of Hawking radiation during evaporation
- We try to look at Hawking radiation as tunneling from supertranslated dynamical Vaidya black hole

# 1. Supertranslation of Vaidya spacetime

□ Vaidya metric in advanced Bondi coordinates

$$\bar{g}_{\mu\nu} dx^\mu dx^\nu = -V dv^2 + 2dvdr + r^2 \gamma_{AB} d\Theta^A d\Theta^B, \quad V \equiv 1 - \frac{2M(v)}{r}$$

□ Supertranslated Vaidya metric:  $g_{\mu\nu} = \bar{g}_{\mu\nu} + \mathcal{L}_\zeta \bar{g}_{\mu\nu}$

$$g_{\mu\nu} dx^\mu dx^\nu = - \left( V - \frac{2fM'}{r} - \frac{MD^2 f}{r^2} \right) dv^2 + 2dvdr - D_A(2Vf + D^2 f) dv d\Theta^A \\ + (r^2 \gamma_{AB} + 2r D_A D_B f - r \gamma_{AB} D^2 f) d\Theta^A d\Theta^B \quad (M' = \partial_v M(v))$$

• We work in the linearized theory in  $f$

✱ The supertranslated metric can be extended from infinity to the interior (BH horizon) since it solves the linearized Einstein eqs. for all  $r$ .


\* Supertranslated Vaidya spacetime contains black hole defined by the trapping horizon

$$r = r_h = 2M + 2fM' + \frac{1}{2}D^2 f$$

\* We can reproduce the supertranslation by injecting a shock wave like  $T_{\mu\nu}$  which has angular dependence

Schwarzschild supertranslation

[Hawking et al.]



# Surface gravity

## □ Surface gravity in spherical symmetry from Kodama vector

$$\nabla^\mu (G_{\mu\nu} K^\mu) = 0, \quad \nabla_\mu K^\mu = 0 \quad [\text{Kodama}]$$

Kodama vector gives a preferred time direction in dynamical situation

- Surface gravity:  $K^\mu \nabla_{[\nu} K_{\mu]} = -\kappa K_\nu$  on horizon [Hayward]

$$\text{For Vaidya BH: } K^\mu = \delta^\mu_v \quad \kappa = (4M(v))^{-1}$$

## □ Extension to supertranslated Vaidya black hole

We find a Kodama-like vector and the surface gravity on the trapping horizon

$$K^\mu = \delta^\mu_v \quad \kappa = \frac{1}{4M} \left( 1 - f \frac{M'}{M} \right)$$

\*Surface gravity has angular dependence due to the supertranslations

## 2. Hawking radiation as tunneling from supertranslated Vaidya BH

□ Hamilton-Jacobi method [Angheben et al., Srinivasan et al.]

(1) KG equation  $\square\phi = 0$  WKB ansatz:  $\phi = A(x)e^{iS/\hbar} + \mathcal{O}(\hbar)$


lowest order:  $g^{\mu\nu}\partial_\mu S\partial_\nu S = 0$  Hamilton-Jacobi equation

(2) Reconstruction of the phase (particle action):

$$S = \int_P dx^\mu \partial_\mu S \quad * \text{Im}S = \text{signal of tunneling}$$

(3) Semiclassical emission rate:

$$\begin{aligned} \Gamma_{\text{em}} &\propto \exp\left(-2\text{Im}S\right) \\ &= e^{-\omega/T} \end{aligned}$$



tunneling path  $\vec{ab}$   
(classically forbidden)

[Hartle et al, Vanzo et al]



□ Supertranslated Vaidya black hole

$$2\partial_r S \partial_v S + \left( V - \frac{2fM'}{r} - \frac{MD^2 f}{r^2} \right) (\partial_r S)^2 \quad \text{Hamilton-Jacobi eq.}$$

$$+ \frac{1}{r^2} D^A (2Vf + D^2 f) \partial_r S \partial_A S + \frac{1}{r^4} (r^2 \gamma^{AB} - 2r D^A D^B f + r \gamma^{AB} D^2 f) \partial_A S \partial_B S = 0$$

- Particle energy:  $\omega \equiv -K^\mu \partial_\mu S = -\partial_v S$   $K^\mu$  : Kodama-like vector
- We concentrate on radial null geodesics:  $\Theta^A = \text{const.} \Rightarrow \partial_A S = 0$

$$\partial_r S_{out} = 2\omega \left( V - \frac{2fM'}{r} - \frac{MD^2 f}{r^2} \right)^{-1}, \quad \partial_r S_{in} = 0$$

$$\Rightarrow S_{out} = - \int \omega dv + \int \frac{2\omega r^2 dr}{Vr^2 - 2fM'r - MD^2 f}, \quad S_{in} = - \int \omega dv.$$

$$\text{Im} S_{out} = \text{Im} \int_{\text{I}} \frac{2\omega r^2 dr}{(r - r_h)(r - r_f)} = \pi\omega \cdot 4M \left( 1 + f \frac{M'}{M} \right)$$


Here  $r = r_h = 2M + 2fM' + \frac{1}{2}D^2 f$  and  $r_f = -\frac{1}{2}D^2 f$



\* Tunneling occurs at the trapping horizon

## □ Semiclassical emission rate

$$\Gamma_{\text{em}} \propto e^{-2\text{Im}S_{\text{out}}} = \exp\left(-8\pi\omega M\left(1 + f\frac{M'}{M}\right)\right) = \exp\left(-\frac{2\pi\omega}{\kappa}\right) = e^{-\omega/T}$$


$$\kappa = \frac{1}{4M} \left(1 - f\frac{M'}{M}\right)$$

\*The emission rate of Hawking radiation has a dependence on supertranslation  $f(\Theta)$ !

\*We only see local temperature of Hawking radiation

## 4. Summary

- Hawking radiation will actually have a dependence on supertranslation  $f(\Theta)$  which disappears in the static limit
- It seems to be possible to distinguish black holes with different supertranslations hair by Hawking radiation
- The semiclassical emission rate can be expressed in terms of the horizon surface gravity  $\kappa$  which can serve as local measure of Hawking temperature

## □ Questions

- Energy flux of Hawking radiation will produce the gravitational memory effects near future null infinity?
- Like energy conservation, Hawking radiation carries a part of charges associated with supertranslation/superrotation? If so, how much amount of the charges?
- What's the implication to information problem?  
[HPS, Bousso etal, Gomez etal, Mirababayi etal, Carney etal, ...]







Backup



## □ BMS supertranslation


- Minkowski space in advanced Bondi coordinates  $(v, r, z, \bar{z})$

$$ds^2 = -dv^2 + 2dvdr + r^2\gamma_{AB}d\Theta^A d\Theta^B$$

$$v = t + r \quad (A, B) = (z, \bar{z})$$

$$z = e^{i\phi} \tan \theta/2$$

$$\gamma_{z\bar{z}} = \frac{2}{(1 + |z|^2)^2} \quad \gamma_{zz} = 0$$



## Asymptotically Minkowski metric

- Bondi gauge:  $g_{rr} = 0, \quad g_{rA} = 0, \quad \partial_r \det \left( \frac{g_{AB}}{r^2} \right) = 0.$

- Boundary conditions at large  $r$  :

$$g_{vv} = -1 + \mathcal{O}\left(\frac{1}{r}\right), \quad g_{vA} = \mathcal{O}(1), \quad g_{vr} = 1 + \mathcal{O}\left(\frac{1}{r^2}\right),$$

$$g_{z\bar{z}} = r^2 \gamma_{z\bar{z}} + \mathcal{O}(1), \quad g_{zz} = \mathcal{O}(r)$$

- Asymptotic expansion of metric  $r \rightarrow \infty$  : past null infinity  $\mathcal{I}^-$

$$ds^2 = -dv^2 + 2dvdr + r^2 \gamma_{AB} d\Theta^A d\Theta^B$$

$$+ \frac{2m}{r} dv^2 + r C_{AB} d\Theta^A d\Theta^B - \left( D^B C_{AB} + \frac{4}{3r} N_A \right) dv d\Theta^A + \dots$$

$$m, C_{AB}, N_A : \text{functions of } (v, z, \bar{z}), \quad \gamma^{AB} C_{AB} = 0,$$

## Infinite degeneracy of BMS vacua

Under the supertranslation, asymptotic data of metric change

$$\delta_f m = f \partial_u m + \frac{1}{4} N^{AB} D_A D_B f + \frac{1}{2} D_A f D_B N^{AB} \quad N_{AB} = \partial_u C_{AB}$$

$$\delta_f C_{AB} = f \partial_u C_{AB} - 2 D_A D_B f + \gamma_{AB} D^2 f$$

- BMS vacuum:  $\partial_u m = 0$ ,  $N_{AB} = 0$ ,

$$D_{\bar{z}}^2 C_{zz} - D_z^2 C_{\bar{z}\bar{z}} = 0 \Rightarrow C_{zz} = -2 D_z^2 C(z, \bar{z})$$

$$\text{But } \delta_f C_{zz} = -2 D_z^2 f \Rightarrow C(z, \bar{z}) \rightarrow C(z, \bar{z}) + f(z, \bar{z})$$

BMS vacua are related by supertranslation.

- Once a vacuum is chosen, supertranslation invariance of BMS vacua is spontaneously broken.
- Different BMS vacua have different angular momentum

# Gravitational memory effect

[Zeldovich et al., Christodoulou, Braginsky et al.]

Before radiation: vacuum with  $C_{AB} = 0$

$$\delta s(z_1, z_2) = L = \sqrt{2} r_0 \gamma_{z\bar{z}} \sqrt{|\delta z|^2}$$

$$\delta z = z_1 - z_2$$


After radiation: vacuum with  $\Delta C_{AB} \neq 0$

Especially  $\Delta C \neq 0$

$$L \rightarrow L + \Delta L \quad \Delta L = \frac{r_0}{2L} (\Delta C_{zz} \delta z^2 + c.c)$$

$$D_z^2 \Delta C_{zz} = 2 \int_{u_i}^{u_f} du \lim_{r \rightarrow \infty} [r^2 T_{uu}^{M+G}] + 2\Delta m$$

(Formular for  $\Delta C$  : Strominger, Zhiboedov)



- $\Delta C$  contains the information of massless/massive matters
- Memory effect: permanent displacement of detectors due to a passage of gravitational wave/null matter  $T_{uu}^{M+G}$  (null memory) and/or a change in state of massive matter  $\Delta m$  (ordinary memory)

## Trapping horizon

Let  $\theta_-$  and  $\theta_+$  be the expansions of the bundles of ingoing and outgoing radial null geodesics, respectively. In our case,

$$\theta_- = -\frac{2}{r}, \quad \theta_+ = \frac{1}{r} \left( V - \frac{2fM'}{r} - \frac{MD^2f}{r^2} \right), \quad \begin{array}{l} \theta > 0 \quad : \text{bundle is expanding} \\ \theta < 0 \quad : \text{bundle is contracting} \end{array}$$

### □ Future outer marginally trapped surface

$$\theta_+ = 0 \quad (\text{marginally trapped}), \quad \theta_- < 0 \quad (\text{future type}), \quad l^\mu \partial_\mu \theta_+ < 0 \quad (\text{outer type})$$

$$\bullet \quad \theta_+ = 0 \Rightarrow r = r_h = 2M + 2fM' + \frac{1}{2}D^2f \quad * \text{Location of horizon}$$

### □ Future outer trapping horizon (FOTH) [Hayward]

- A foliation of future outer marginally trapped surface

\* FOTH gives the local definition of supertranslated Vaidya black hole

## □ Tunneling in Schwarzschild black hole

HJ eq.:  $\left(1 - \frac{2M}{r}\right)(\partial_r S)^2 - 2\omega\partial_r S = 0$  Trajectory on  $(v - r)$  plane

$$\Rightarrow \partial_r S_{out} = \frac{2\omega}{1 - \frac{2M}{r}}, \quad \partial_r S_{in} = 0$$


• particle energy:  $\omega = -\xi^\mu p_\mu = -\xi^\mu \partial_\mu S = -\partial_v S$

Particle action is given by a line integral along a path

$$S_{out} = - \int \omega dv + \int \frac{2\omega dr}{1 - \frac{2M}{r}}$$

Imaginary part of the action is

$$\begin{aligned} \text{Im} S_{out} &= \text{Im} \int_{a \rightarrow b \rightarrow c} \frac{2\omega dr}{1 - \frac{2M}{r}} = \text{Im} \int_{a \rightarrow b} \frac{2\omega dr}{1 - \frac{2M}{r}} \\ &= \text{Im} \int_{a \rightarrow b} \frac{2\omega r dr}{r - 2M - i\epsilon} = 4\pi\omega M. \end{aligned}$$




## Semiclassical emission rate

$$\Gamma_{\text{em}} \propto \exp(-2\text{Im}S_{\text{out}}) = \exp(-8\pi\omega M) = \exp(-\omega/T_{\text{H}})$$

We can read off the Hawking temperature:  $T_{\text{H}} = \frac{\kappa_s}{2\pi}$ ,  $\kappa_s = \frac{1}{4M}$

# X □ Two types of tunneling path for $S_{out}$



- Type-I: classically forbidden trajectory backward in time
- Type-II: classically allowed trajectory forward in time.

[Vanzo et al.]

Along these paths,

$$v = \text{const.} \quad (\text{type-I}),$$

$$\frac{dr}{dv} = \frac{1}{2} \left( V - \frac{2fM'}{r} - \frac{MD^2f}{r^2} \right) \quad (\text{type-II}).$$

For type-II path, we have

$$S_{out} = - \int_{\text{II}} dv \omega + \int_{\text{II}} \partial_r S_{out} dr = 0$$

No tunneling at all.



## X □ Entropy and semiclassical emission rate

$$S_{\text{dyn}} = \frac{A_h}{4} = 4\pi M^2 + \sqrt{16\pi a_{00}} M M'$$

$$f(\Theta) = \sum_{l,m} a_{lm} Y_{lm}(\Theta)$$

After an emission of Hawking radiation,  $M \rightarrow M - \omega$ ,  $S_{\text{dyn}} \rightarrow S_{\text{dyn}} + \Delta S$

$$\Delta S = -\omega(8\pi M + \sqrt{16\pi a_{00}} M') + \mathcal{O}(\omega^2)$$

$$\text{Thus, } \Delta S = \int \frac{d\Omega}{4\pi} \ln \Gamma_{\text{em}}$$

$$\Rightarrow \frac{d\Delta S}{d\Omega} = \ln \Gamma_{\text{em}} \quad \text{Hawking radiation carries away from the black hole different amount of entropy at different angles}$$