

A Few Thoughts on Gravity and Cosmology

- geemunu, geemunu, geemunu...-

Misao Sasaki
YITP, Kyoto University

a bit of recollections

cosmological perturbation
theory (CPT)

collaboration with
H Kodama

gravitational waves
(GW)

collaboration with
T Futamase through
TAMA300

(+ gravitational lensing)

Progress of Theoretical Physics Supplement No. 78, 1984

Cosmological Perturbation Theory

Hideo KODAMA and Misao SASAKI*

Department of Physics, University of Tokyo, Tokyo 11

**Department of Physics, Kyoto University, Kyoto 606*

(Received September 5, 1984)

The linear perturbation theory of spatially homogeneous and isotropic universe is reformulated extensively. In the first half of the article, a gauge-invariant formalism for the theory is carried out with special attention paid to the geometrical meaning of the perturbation. In the second half of the article, the application of the theory to some important models is discussed.

PHYSICAL REVIEW D

VOLUME 40, NUMBER 8

15 OCTOBER 1989

Light propagation and the distance-redshift relation in a realistic inhomogeneous universe

Toshifumi Futamase

Department of Physics, Faculty of Science, Hirosaki University, Hirosaki 036, Japan

Misao Sasaki

*NASA/Fermilab Astrophysics Center, Fermi National Accelerator Laboratory, Batavia, Illinois 60510
and Research Institute for Theoretical Physics, Hiroshima University, Takehara, Hiroshima 725, Japan**

(Received 3 May 1989)

We investigate the propagation of light rays in a clumpy universe constructed by a cosmological version of the post-Newtonian approximation. We show that the linear approximation to the propagation equations is valid in the region $z \lesssim 1$ even if the density contrast is much larger than unity. Based on a general order-of-magnitude statistical consideration, we argue that the linear approximation is still valid for $z \gtrsim 1$. Then we give a general formula for the distance-redshift relation in a clumpy universe and derive an explicit expression for a simplified situation in which the effect of the gravitational potential of inhomogeneities dominates. In the light of the derived relation we discuss the validity of the Dyer-Roeder distance. Furthermore, we consider a simple model of an inhomogeneous universe and investigate statistical properties of light rays. We find that the result of this specific example also supports the validity of the linear approximation.

a bit of recollections

cosmological perturbation
theory (CPT)

gravitational waves
(GW)

- during 80's, GWs were regarded as more realistic, of firm GR foundation.
- during 90's, CPT became realistic, thanks to COBE/DMR measured anisotropy.
- during 00's, both became realistic. But...

fairy tales are necessary for healthy growth of **children**
(H Sato at a theory group workshop, 中間発表会, in '90s)

(chukan-happyo-kai)

so **WE** (at least **I**) need fairy tales...

Gravity Today

- No deviation from **General Relativity**

cf. Will's living review '06

- **solar system tests – PPN parameters**

$$g_{00} = -1 + 2\psi - 2\beta\psi^2 + \dots : \beta_{GR} = 1$$

$$g_{ij} = \delta_{ij} (1 + 2\gamma\psi + \dots) : \gamma_{GR} = 1$$

$|\gamma - 1| < 2.3 \times 10^{-5}$: Shapiro time delay (Bertotti et al. '03)

$|4\beta - \gamma - 3| < 4.4 \times 10^{-4}$: Strong EP (Baessler et al. '99)

- **constancy of gravitational constant**

$|d \log G / dt| < 10^{-12} \text{ yr}^{-1}$: Lunar laser ranging (Williams et al '04)

- binary pulsar – GW emission rate


Hulse-Taylor binary (B1913+16)

- orbital change due to GW emission

$$\frac{\dot{P}_{\text{B1913+16}}}{\dot{P}_{\text{GR}}} = 1.0013 \pm 0.0021$$

perfect match with
GR predictions


- periastron time shift



gravitational wave astronomy will further test GR


Cosmology Today

- **Big Bang** theory has been firmly established



another strong evidence for **GR**

Strong evidence for Inflation



$$\langle \Delta T(\vec{n}_1) \Delta T(\vec{n}_2) \rangle = \sum_{\ell=0}^{\infty} \frac{2\ell+1}{4\pi} C_{\ell} P_{\ell}(\cos \theta).$$

- highly Gaussian fluctuations
- almost scale-invariant spectrum

only to be confirmed (by tensor modes?)

Fundamental(?) Issues

- **Dark Matter**

Is it really `matter`?

Perhaps yes, because it **gravitates**.

fermion? boson? primordial BH?
something else?

Is there a way to **generically distinguish them?**

- **Dark Energy**

- **apparent** accelerated expansion of the universe

Is the expansion really accelerating?

e.g. inhomogeneous universe models

How can we **confirm acceleration**?

- **modified gravity** vs **unknown matter field**

How to distinguish?

large scale structure formation

$w < -1$ implies modified gravity, etc...

Can we **falsify GR**?

any other **effective discriminators**?

comments on modified gravity



- **time** is important in spacetime!
- any smooth function $f(x)$ contains infinitely large Fourier modes...
- amplitude of Riemann? in which frame?

- **Inflation**

- How did inflation begin?

- what guarantees homogeneity and isotropy?
quantum cosmology/gravity?

- What is 'inflaton'?

- what determines the end of inflation?

- flatness / open inflation?

- non-Gaussianity? tensor-scalar ratio?

- eternal inflation, anthropic principle,
probability measure,...**

new guiding principle / working hypothesis?

What's next?
Which direction?



With a bit of 我田引水 (ga den-in sui)
which means 'self advocacy', more or less...

String Theory Landscape!


Isn't there a landscape in LQG?
(a question to Abhay)



String theory landscape

Bousso & Pochinski ('00), Susskind, Douglas, KKLT ('03), ...

- There are $\sim 10^{500}$ vacua in string theory
 - vacuum energy ρ_v may be positive or negative
 - typical energy scale $\sim M_P^4$
 - some of them have $\rho_v \ll M_P^4$




testing string theory landscape in cosmology?



Cosmic Landscape

various vacua realized in the early universe



distribution determined by various factors

probability measure, density of states,
quantum equilibrium, ...

quantum transitions between various vacua

Creation of open universe


de Sitter (dS) vacuum: $O(4,1)$



Euclidean bubble: $O(4)$



nucleated bubble: $O(3,1)$
= open universe



Our Universe was born out of quantum tunneling!

➤ two possibilities

1. inflation after tunneling was short enough ($N \sim 60$)

$$1 - \Omega_0 = 10^{-2} \sim 10^{-3} \quad \text{“open universe”}$$

➔ signatures in large angle CMB anisotropies

Yamauchi, Linde, Naruko, Tanaka & MS (2011)

2. inflation after tunneling was long enough ($N \gg 60$)


$$1 - \Omega_0 \ll 1 \quad \text{“flat universe”}$$

➔ signatures from bubble collisions

Sugimura, Yamauchi & MS (2012)


Sugimura's talk:
parallel session
on Monday

trace of bubble collisions



➤ simple model

- no (spherically symmetric) bubble seen in the CMB map
 - ⇔ negligible effect on curvature perturbation (\sim Newton potential) at leading order



bubbles may be seen as “localized” non-Gaussianity


$$\Phi(\mathbf{x}) = \Phi_{Gauss}(\mathbf{x}) + \underbrace{f_{NL}(\mathbf{x})}_{\text{space-dependent}} \Phi_{Gauss}^2(\mathbf{x}) + \dots$$



space-dependent

Non-Gaussian bubbles in the CMB Sky

Sugimura, Yamauchi & MS (2012)



detection of a **spherically symmetric “localized” non-Gaussianity** will be the first observational **signature of string theory!**

Summary without summary...

We are entering an era of
precision cosmology
gravitational wave astronomy

any tiny deviation from GR would be revolutionary

develop 'realistic' GR cosmology

perturbative, non-perturbative, numerical, observational...

continued to T Futamase's talk